

# New Combinatorial Characterizations of Generalized Quadrangles

J. A. THAS

## 1. INTRODUCTION

Let  $\mathcal{S} = (\mathcal{P}, \mathcal{B}, I)$  be a generalized quadrangle of order  $(s, t)$ . For a fixed point  $p$ , define the following condition:  $(M)_p$  For any two lines  $L, M$  of  $\mathcal{S}$  incident with  $p$ , the group of collineations of  $\mathcal{S}$  fixing  $L$  and  $M$  pointwise and  $p$  linewise is transitive on the lines ( $\neq L$ ) incident with a given point  $x$  on  $L$  ( $x \neq p$ ).

Then  $\mathcal{S}$  is said to satisfy condition  $(M)$  provided it satisfies  $(M)_p$  for all points  $p \in \mathcal{P}$ . Let  $(\hat{M})$  be the dual of  $(M)$ . If  $\mathcal{S}$  satisfies both  $(M)$  and  $(\hat{M})$  and if  $s \neq 1 \neq t$ , it is called a *Moufang generalized quadrangle* [12], and by a celebrated theorem of Tits [11, 13] a *thick generalized quadrangle is a Moufang generalized quadrangle iff it is one of the classical examples or their duals*. Recently Tits proved that the theorem is also valid in the infinite case [13, 14].

Another characterization of all classical generalized quadrangles was given by Buekenhout and Lefèvre [2]: *The generalized quadrangle  $\mathcal{S}$  of order  $(s, t)$  is classical iff it is embeddable in a projective space  $PG(n, s)$* . In the infinite case the theorem was proved by K. J. Dienst [4].

Here we introduce a new condition  $(A)$ , and its dual  $(\hat{A})$ , for generalized quadrangles (Section 3), and we show (Section 4) *that a thick generalized quadrangle is a classical or a dual classical generalized quadrangle iff it satisfies one of the conditions  $(A)$  or  $(\hat{A})$* . Finally, in Section 5 a new characterization of the dual of  $T(\mathcal{O})$ , the Tits' quadrangle arising from an ovoid  $\mathcal{O}$ , is given.

## 2. BASIC CONCEPTS AND RESULTS

A generalized quadrangle of order  $(s, t)$  ( $s \geq 1, t \geq 1$ ) is an incidence structure  $\mathcal{S} = (\mathcal{P}, \mathcal{B}, I)$  with pointset  $\mathcal{P}$ , lineset  $\mathcal{B}$ , and symmetric point–line incidence relation  $I$  satisfying the following axioms:

- (i) *Each point (resp. line) is incident with  $1+t$  lines (resp.  $1+s$  points).*
- (ii) *Two points are incident with at most one line.*
- (iii) *If  $x$  is a point not incident with a line  $L$ , there is a unique point–line pair  $(y, M) \in \mathcal{P} \times \mathcal{B}$  with  $xIMyIL$ .*

If  $s \neq 1 \neq t$ ,  $\mathcal{S}$  is called *thick*. When the points  $x, y$  (resp. lines  $L, M$ ) of  $\mathcal{S}$  are collinear (resp. concurrent), we write  $x \sim y$  (resp.  $L \sim M$ ); otherwise  $x \not\sim y$  (resp.  $L \not\sim M$ ). For  $(x, L) \in \mathcal{P} \times \mathcal{B}$  put  $x^\perp = \{y \in \mathcal{P} \mid y \sim x\}$  and  $L^\perp = \{M \in \mathcal{B} \mid L \sim M\}$ . The *trace* of a pair of distinct points  $(x, y)$  is defined to be the set  $x^\perp \cap y^\perp$  and is denoted  $\{x, y\}^\perp$ . More generally, if  $A \subset \mathcal{P}$ , a “perp.” is defined by  $A^\perp = \cap \{x^\perp \mid x \in A\}$ . For  $x \neq y$ , the *span* of the pair  $(x, y)$  is  $\{x, y\}^{\perp\perp} = \{u \in \mathcal{P} \mid u \in z^\perp \text{ for all } z \in x^\perp \cap y^\perp\}$ . And the *closure* of the pair  $(x, y)$  is  $\text{cl}(x, y) = \{z \in \mathcal{P} \mid z^\perp \cap \{x, y\}^{\perp\perp} \neq \emptyset\}$ . We say that the pair  $(x, y)$  is *regular* if either  $x \sim y$  or  $x \not\sim y$  and  $|\{x, y\}^{\perp\perp}| = t+1$ . The point  $x$  is *regular* provided  $(x, y)$  is regular for all  $y \in \mathcal{P}, y \neq x$ . If  $\mathcal{S}$  contains a regular pair of non-collinear points, then  $s = 1$  or  $s \geq t$ . A *triad (of points)* is a triple of pairwise non-collinear points.

There is a point–line duality for generalized quadrangles (of order  $(s, t)$ ) which interchanges “point” and “line”, interchanges  $s$  and  $t$ , in any definition or theorem. Normally,

we assume without further notice that the dual of a given theorem or definition is also given. A line  $L$  is *coregular* provided each point incident with  $L$  is regular.

The generalized quadrangle arising from a non-singular hyperquadric  $Q$  of index 2 in  $PG(d, q)$ ,  $d = 3, 4$  or  $5$ , is denoted by  $Q(d, q)$ ; the generalized quadrangle arising from a non-singular Hermitian variety  $H$  in  $PG(d, q^2)$ ,  $d = 3$  or  $4$ , is denoted by  $H(d, q^2)$ ; the generalized quadrangle formed by the points of  $PG(3, q)$  together with all totally isotropic lines with respect to a symplectic polarity of  $PG(3, q)$  is denoted by  $W(3, q)$ . These are the so-called *classical* generalized quadrangles. We remark that  $W(3, q)$  is isomorphic to the dual of  $Q(4, q)$ , that  $H(3, q^2)$  is isomorphic to the dual of  $Q(5, q)$ , and that all classical generalized quadrangles but  $Q(3, q)$  are thick [10].

The Tits' quadrangle arising from an ovoid  $\mathcal{O}$  of  $PG(3, q)$  [3] will be denoted by  $T(\mathcal{O})$ . It is isomorphic to  $Q(5, q)$  iff  $\mathcal{O}$  is an elliptic quadric [10].

The generalized quadrangle  $\mathcal{S}$  is said to satisfy *property (H)* provided  $x \in \text{cl}(y, z)$  implies  $y \in \text{cl}(x, z)$  for any triad  $(x, y, z)$ . Also we introduce the following *axiom ( $\hat{D}$ )*: If  $l_1, l_2, m_1, m_2, u_1, u_2, n_1$  are distinct points for which  $l_1 \not\sim m_2, l_1 \sim l_2, m_1 \sim m_2, l_1 \sim u_1 \sim m_1, l_2 \sim u_2 \sim m_2, u_1 \sim u_2, l_1 \sim n_1 \sim m_1$  and for which  $l_1 l_2, m_1 m_2, l_1 u_1, u_1 m_1, l_2 u_2, u_2 m_2, u_1 u_2, l_1 n_1, n_1 m_1$  are distinct lines, then there exists a point  $n_2$  with  $l_2 \sim n_2 \sim m_2$  and  $n_1 \sim n_2$ . Theorem (a) and the dual of (b) are proved in [10, 8] and [9] respectively.

(a) If  $\mathcal{S}$  satisfies property (H), then one of the following must occur:

- (i) each point of  $\mathcal{S}$  is regular (implying  $s = 1$  or  $s \geq t$ ),
- (ii)  $|\{x, y\}^{\perp\perp}| = 2$  for all  $x, y \in \mathcal{P}$  with  $x \not\sim y$ , or
- (iii)  $\mathcal{S}$  is isomorphic to  $H(4, s)$ .

(b) If  $2 < t < s$  and  $\mathcal{S}$  contains a coregular line  $L$  such that ( $\hat{D}$ ) is satisfied whenever  $l_1$  or  $l_2$  is on  $L$ , then  $s = t^2$  and  $\mathcal{S}$  is isomorphic to the dual of a  $T(\mathcal{O})$  of Tits.

If  $\mathcal{S}$  is a generalized quadrangle with parameters  $s, t$ , where  $2 < t < s$ , for which each point is regular and for which ( $\hat{D}$ ) is satisfied, then  $t^2 = s$  and  $\mathcal{S}$  is isomorphic to  $H(3, s)$ .

From (b) there easily follows

(c) If  $2 < t < s$  and  $\mathcal{S}$  contains a coregular line  $L$  such that ( $\hat{D}$ ) is satisfied whenever  $l_1$  (resp.  $l_2$ ) is on  $L$ , then  $s = t^2$  and  $\mathcal{S}$  is isomorphic to the dual of a  $T(\mathcal{O})$  of Tits.

**PROOF OF (c).** Suppose that ( $\hat{D}$ ) is satisfied whenever  $l_1$  is on the coregular line  $L$ . Then we prove that ( $\hat{D}$ ) holds whenever  $l_2$  is on  $L$ . Since  $l_2$  is on  $L$ , the point  $l_2$  is regular. If  $l_2 \sim m_1$ , then  $u_2, m_1, l_1 \in \{l_2, u_1\}^{\perp}$ , and so  $m_2 \sim l_1$ , a contradiction. So we always have  $l_2 \not\sim m_1$ . Let  $v_2$  be a point for which  $v_2 \sim l_2, v_2 \sim m_2$  and  $v_2 \neq u_2$ . Then, by axiom ( $\hat{D}$ ), there is a point  $v_1$  such that  $v_1 \sim l_1, v_1 \sim m_1, v_1 \sim v_2$ . Since  $l_2 \not\sim m_1$  we have  $v_1 \neq u_1$ . From  $l_2 \not\sim m_1$  it also follows that the point  $v_1$  is uniquely defined by  $v_2$ , and that  $v_2 \mapsto v_1$  defines an injection from  $\{l_2, m_2\}^{\perp} - \{u_2\}$  into  $\{l_1, m_1\}^{\perp} - \{u_1\}$ . Since  $|\{l_2, m_2\}^{\perp} - \{u_2\}| = |\{l_1, m_1\}^{\perp} - \{u_1\}| = t$ , the injection  $v_2 \mapsto v_1$  is a bijection from  $\{l_2, m_2\}^{\perp} - \{u_2\}$  onto  $\{l_1, m_1\}^{\perp} - \{u_1\}$ . It immediately follows that ( $\hat{D}$ ) is satisfied whenever  $l_2$  is on  $L$ . And so, the first part of Theorem (b) applies.

The other case is completely analogous.

### 3. PROPERTIES (A) AND ( $\hat{A}$ )

Let  $\mathcal{S} = (\mathcal{P}, \mathcal{B}, I)$  be a generalized quadrangle of order  $(s, t)$ . If  $\mathcal{B}^{\perp\perp} = \{\{x, y\}^{\perp\perp}, x \not\sim y\}$ , then let  $\mathcal{S}^{\perp\perp} = (\mathcal{P}, \mathcal{B}^{\perp\perp}, \in)$ . For  $x \in \mathcal{P}$  we say  $\mathcal{S}$  satisfies *property (A)<sub>x</sub>* if for any  $M = \{y, z\}^{\perp\perp} \in \mathcal{B}^{\perp\perp}$  with  $x \in \{y, z\}^{\perp}$ , and any  $u \in \text{cl}(y, z) \cap (x^{\perp} - \{x\})$  (or  $u \neq x$  and  $u \perp x$  for some  $k \in M$ ) with  $u \notin M$ , the substructure of  $\mathcal{S}^{\perp\perp}$  generated by  $M$  and  $u$  is a dual affine plane. The generalized quadrangle  $\mathcal{S}$  is said to satisfy *property (A)* if it satisfies (A)<sub>x</sub> for

all  $x \in \mathcal{P}$ . The duals of these properties are denoted resp.  $(\hat{A})_L$  and  $(\hat{A})$ . We notice that property (A) is equivalent to the following: if  $M = \{y, z\}^{\perp\perp} \in \mathcal{B}^{\perp\perp}$  and  $u \in \text{cl}(y, z)$ ,  $u \notin M \cup M^\perp$ , then  $u$  and  $M$  generate a dual affine plane in  $\mathcal{S}^{\perp\perp}$ .

#### 4. THEOREM

*The thick generalized quadrangle  $\mathcal{S}$  is a classical or a dual classical generalized quadrangle iff it satisfies one of the conditions (A) or  $(\hat{A})$ .*

PROOF. It is not difficult (and an interesting exercise) to check that a thick classical or dual classical generalized quadrangle satisfies one of the conditions (A) or  $(\hat{A})$ .

Next let us assume that the thick generalized quadrangle  $\mathcal{S} = (\mathcal{P}, \mathcal{B}, I)$  satisfies (A). We shall prove that property (H) is satisfied. So we consider a triad  $(u, y, z)$  for which  $u \in \text{cl}(y, z)$ . Let  $\pi$  be the dual affine plane generated by  $\{y, z\}^{\perp\perp}$  and  $u$  in  $\mathcal{S}^{\perp\perp}$ . Evidently  $\{z, u\}^{\perp\perp}$  is a line of  $\pi$ . In  $\pi$  the point  $y$  is not collinear with exactly one point of  $\{z, u\}^{\perp\perp}$ , i.e., in  $\mathcal{S}$  the point  $y$  is collinear with exactly one point of  $\{z, u\}^{\perp\perp}$ . Hence  $y \in \text{cl}(z, u)$ , and so (H) is satisfied.

Now we assume that  $\mathcal{S} \not\cong H(4, s)$ . If  $|\{y, z\}^{\perp\perp}| = 2$  for all  $y, z \in \mathcal{P}$  with  $y \not\sim z$ , then for any  $M = \{y, z\}^{\perp\perp} \in \mathcal{B}^{\perp\perp}$  and any  $u \in \text{cl}(y, z) - \{y, z\}^\perp$  with  $u \notin M$  (such a  $u$  exists by the thickness of  $\mathcal{S}$ ), the substructure of  $\mathcal{S}^{\perp\perp}$  generated by  $M$  and  $u$  has three points and consequently is not a dual affine plane, a contradiction. Then, from Section 2, it follows that every point of  $\mathcal{S}$  is regular. Since  $\mathcal{S}$  is thick we have  $t \neq 1$  and  $s \geq t$ . If  $s = t$ , then it is well known that  $\mathcal{S}$  is isomorphic to  $W(3, s)$  [10]. If  $t = 2 < s$ , then  $\mathcal{S}$  is isomorphic to  $H(3, 4)$  [5, 6, 7]. Hence we now suppose that  $2 < t < s$ .

We shall prove that  $\mathcal{S}$  satisfies property  $(\hat{D})$ . Consider points  $l_1, l_2, m_1, m_2, u_1, u_2, n_1$  as in axiom  $(\hat{D})$ . Since the pair  $(l_2, m_2)$  is regular the point  $u_1$  is element of  $\text{cl}(l_2, m_2)$ . We consider the dual affine plane  $\pi$  generated by  $\{l_2, m_2\}^{\perp\perp}$  and  $u_1$  in  $\mathcal{S}^{\perp\perp}$ . Lines of  $\pi$  are  $\{l_2, u_1\}^{\perp\perp}$  and  $\{m_2, u_1\}^{\perp\perp}$ . Since  $l_1$  (resp.  $m_1$ ) is an element of  $\{u_1, l_2\}^\perp$  (resp.  $\{u_1, m_2\}^\perp$ ), the line  $l_1 n_1$  (resp.  $m_1 n_1$ ) contains a point  $r$  (resp.  $r'$ ) of  $\{u_1, l_2\}^{\perp\perp}$  (resp.  $\{u_1, m_2\}^{\perp\perp}$ ). Then  $R = \{r, r'\}^{\perp\perp}$  is a line of  $\pi$ . As any two lines of  $\pi$  intersect, the lines  $\{l_2, m_2\}^{\perp\perp}$  and  $R$  of  $\pi$  have a point  $r''$  in common. We have  $r'' \sim n_1$ . If  $n_2$  is the point of  $n_1 r''$  which is collinear (in  $\mathcal{S}$ ) with  $l_2$ , then  $m_2 \in \{l_2, r''\}^{\perp\perp}$  and  $n_2 \in \{l_2, r''\}^\perp$  imply  $n_2 \sim m_2$ . Hence  $n_2 \sim n_1$ ,  $n_2 \sim l_2$ , and  $n_2 \sim m_2$ . Consequently  $(\hat{D})$  is satisfied, and so  $\mathcal{S}$  is isomorphic to  $H(3, s)$ .

We have proved that if the thick quadrangle  $\mathcal{S}$  satisfies (A), then  $\mathcal{S}$  is isomorphic to  $H(4, s)$ ,  $W(3, s)$ , or  $H(3, s)$ . Hence, if the thick quadrangle  $\mathcal{S}$  satisfies one of the conditions (A) or  $(\hat{A})$ , then it is isomorphic to  $H(4, s)$ , the dual of  $H(4, t)$ ,  $W(3, s)$ ,  $Q(4, s)$ ,  $H(3, s)$  or  $Q(5, s)$ .

#### 5. THEOREM

*Let  $\mathcal{S}$  be a thick generalized quadrangle with  $s \neq t$ . Then  $\mathcal{S}$  is isomorphic to the dual of a Tits' quadrangle  $T(\mathcal{O})$ , arising from an ovoid  $\mathcal{O}$ , iff  $(A)_x$  is satisfied for all points  $x$  incident with some coregular line  $L$ .*

PROOF. Let  $\mathcal{S}$  be the dual of the Tits' quadrangle  $T(\mathcal{O})$  arising from the ovoid  $\mathcal{O}$  of  $PG(3, q)$ . Then property  $(A)_x$  is satisfied for every point  $x$  incident with the (coregular) line of type (iii) [3].

Now let  $\mathcal{S}$  be a thick generalized quadrangle with  $s \neq t$  such that  $(A)_x$  is satisfied for every point  $x$  on some coregular line  $L$ . Since  $\mathcal{S}$  contains regular points and  $1 < s \neq t$ , we have  $s > t > 1$ . We shall prove that  $(\hat{D})$  is satisfied whenever  $l_1$  is on  $L$ . Hence consider points  $l_1, l_2, m_1, m_2, u_1, u_2, n_1$  as in axiom  $(\hat{D})$ . Since the pair  $(l_2, u_1)$  is regular the

point  $n_1$  is an element of  $\text{cl}(l_2, u_1)$ . We consider the dual affine plane  $\pi$  generated by  $\{l_2, u_1\}^{\perp\perp}$  and  $n_1$  in  $\mathcal{S}^{\perp\perp}$ . Lines of  $\pi$  are  $\{u_1, n_1\}^{\perp\perp}$  and  $\{l_2, n_1\}^{\perp\perp}$ . Since  $u_2$  (resp.  $m_1$ ) is an element of  $\{l_2, u_1\}^{\perp}$  (resp.  $\{u_1, n_1\}^{\perp}$ ), the line  $u_2m_2$  (resp.  $m_1m_2$ ) contains a point  $r$  (resp.  $r'$ ) of  $\{l_2, u_1\}^{\perp\perp}$  (resp.  $\{u_1, n_1\}^{\perp\perp}$ ). Then  $R = \{r, r'\}^{\perp\perp}$  is a line of  $\pi$ . As any two lines of  $\pi$  intersect, the lines  $\{l_2, n_1\}^{\perp\perp}$  and  $R$  of  $\pi$  have a point  $r''$  in common. We have  $r'' \sim m_2$ . If  $n_2$  is the point of  $m_2r''$  which is collinear (in  $\mathcal{S}$ ) with  $l_2$ , then from  $n_1 \in \{l_2, r''\}^{\perp\perp}$  and  $n_2 \in \{l_2, r''\}^{\perp}$  it follows that  $n_2 \sim n_1$ . Hence  $n_2 \sim n_1$ ,  $n_2 \sim l_2$ , and  $n_2 \sim m_2$ . Consequently  $\hat{D}$  is satisfied. If  $t \neq 2$ , Theorem (c) in Section 2 tells us that  $s = t^2$  and that  $\mathcal{S}$  is isomorphic to the dual of a Tits' quadrangle  $T(\mathcal{O})$  arising from an ovoid  $\mathcal{O}$ . If  $t = 2$ , then it is well known that  $\mathcal{S}$  is isomorphic to  $H(3, 4)$ , the unique generalized quadrangle of order  $(4, 2)$  [5, 6, 7].

### COROLLARIES

(a) Let  $\mathcal{S}$  be a thick generalized quadrangle with  $s \neq t$  and  $s$  odd. Then  $\mathcal{S}$  is isomorphic to  $H(3, s)$  iff  $(A)_x$  is satisfied for all points  $x$  incident with some coregular line  $L$ .

PROOF. If  $\mathcal{S} \cong H(3, s)$ , then  $(A)_x$  is satisfied for every point  $x$  of  $\mathcal{S}$ . Conversely, if  $(A)_x$  is satisfied for all points  $x$  incident with some coregular line  $L$ , then  $\mathcal{S}$  is isomorphic to the dual of  $T(\mathcal{O})$  for some ovoid  $\mathcal{O}$ . Since  $s$  is odd,  $\mathcal{O}$  is an elliptic quadric [1], and so  $T(\mathcal{O}) \cong Q(5, s)$ . Hence  $\mathcal{S} \cong H(3, s)$ .

(b) Let  $\mathcal{S}$  be a thick generalized quadrangle with  $s \neq t$  and all points regular. Then  $\mathcal{S}$  is isomorphic to  $H(3, s)$  iff  $(A)_x$  is satisfied for all points  $x$  incident with some line  $L$ .

PROOF. If  $\mathcal{S} \cong H(3, s)$ , then  $(A)_x$  is satisfied for every point  $x$  of  $\mathcal{S}$ . Conversely, if  $(A)_x$  is satisfied for all points  $x$  incident with some coregular line  $L$ , then  $\mathcal{S}$  is isomorphic to the dual of  $T(\mathcal{O})$  for some ovoid  $\mathcal{O}$ . Since all points of  $\mathcal{S}$  are regular, all lines of  $T(\mathcal{O})$  are regular, and so all ovals on  $\mathcal{O}$  are conics. Consequently  $\mathcal{O}$  is an elliptic quadratic [1]. Hence  $T(\mathcal{O}) \cong Q(5, s)$  and so  $\mathcal{S} \cong H(3, s)$ .

### REFERENCES

1. A. Barlotti, Some topics in finite geometrical structures, University of North Carolina at Chapel Hill, Institute of Statistics Mimeo Series No. 439, 1965.
2. F. Buekenhout and C. Lefèvre, Generalized quadrangles in projective spaces, *Arch. Math.* **25** (1974), 540–552.
3. P. Dembowski, *Finite Geometries*, Springer-Verlag, Berlin, 1968.
4. K. J. Dienst, Verallgemeinerte Vierecke in Projektiven Räumen, Technische Hochschule Darmstadt, Preprint Nr. 491, 1979.
5. J. J. Seidel, Strongly regular graphs with  $(-1, 1, 0)$  adjacency matrix having eigenvalue 3, *Linear Algebra Appl.* **1** (1968), 281–298.
6. E. Shult, Characterizations of certain classes of graphs, *J. Combin. Theory Ser. B* **13** (1972), 142–167.
7. J. A. Thas, On 4-gonal configurations with parameters  $r = q^2 + 1$  and  $k = q + 1$ , *Geom. Dedicata* **3** (1974), 365–375.
8. J. A. Thas, On generalized quadrangles with parameters  $s = q^2$  and  $t = q^3$ , *Geom. Dedicata* **5** (1976), 485–496.
9. J. A. Thas, Combinatorial characterizations of generalized quadrangles with parameters  $s = q$  and  $t = q^2$ , *Geom. Dedicata* **7** (1978), 223–232.
10. J. A. Thas and S. E. Payne, Classical finite generalized quadrangles: a combinatorial study, *Ars Combin.* **2** (1976), 57–110.
11. J. Tits, Classification of buildings of spherical type and Moufang polygons: a survey, *Atti Conv. Lincei (Coll. Geom. Comb. Roma 1973)* **17** (1976), 230–246.

12. J. Tits, *Buildings and BN-pairs of Spherical Type*, Springer-Verlag Lecture Notes No. 386, Berlin, 1974.
13. J. Tits, *Quadrangles de Moufang, I*, to appear.
14. J. Tits, private communication.

*Received 1 August 1980 and in revised form 6 April 1981*

J. A. THAS  
*Seminar of Geometry and Combinatorics, University of Ghent,  
Krijgslaan 271, 9000 Gent, Belgium*